Novel Chemistry at Extreme Conditions

Maddury Somayazulu, S. G. Gramsch, Muhtar Ahart, Ivan Naumov, Russell J Hemley (*CDAC team*)

Geophysical Laboratory

Carnegie Institution of Washington

Components of the Center

Academic Partners

- CARNEGIE INST. (Hemley)
- CALTECH (Fultz)
- GEORGETOWN (Ichiye)
- NORTHWESTERN UNIV. (Jacobsen)
- UCLA (Kavner)
- UNIV. ALABAMA – BIRMINGHAM (Vohra)
- UNIV. at BUFFALO (Zurek)
- UNIV. CALIF. – BERKELEY (Wenk & Jeanloz)
- UNIV. HAWAI'I (Dera)
- UNIV. ILLINOIS (Dlott & Cahill)
- UNIV. UTAH (Miyagi)
- WASHINGTON UNIV. (Schilling)
- WASHINGTON STATE UNIV. (Yoo)
- YALE UNIV. (Lee)

Academic Collaborators

FACILITY USERS

NNSA Laboratory Partners

ALL HIGH *P-T* GROUPS AT LLNL, LANL, SNL;
STEERING/ADVISORY COMMITTEE MEMBERS

CDAC collaborations and related Research Projects

Muhtar Ahart, Takaki Muramatsu, Tim Strobel, Stephen Gramsch, Ivan Naumov, Hanyu Liu, Ajay Mishra, Alex Goncharov , Sergey Lobanov, Zhenxian Liu, Jerry Potter, George Cody, Ho-Kwang Mao, Larry Finger and Russell Hemley at the **Geophysical Laboratory.**

Raja Chellappa and Dana Dattelbaum from **LANL** – *Hydrazine and Xe-N_2*

Przemek Dera from **GL to GSECARS to Univ. Hawaii** – *Xe-H_2 and Xe-Cl_2*

Choong-Shik Yoo from **WSU** – *Hydride superconductors*

Jennifer Ciezak from **ARL** – *Boron Carbide and doped Boron Carbide*

Eva Zurek from **SUNY** – *Boron Carbide and Alkali Superhydrides*




Beamtime Support and Techniques Collaborators

- Maria Baldini (APS); Chen Li (SNS); Zhenxian Liu (NSLS)
- Viktor Struzhkin, Alex Goncharov and Reini Boehler at GL
- Przemek Dera and Vitali Prakapenka at GSECARS
- Yue Meng, Stas Sinogeikin and Changyong Park at HPCAT
- Peter Liermann at PETRA

Science News

from research organizations

Hydrogen-economy on the way? New hydrogen-storage method discovered

Date: November 25, 2009

Source: Carnegie Institution

Summary: Scientists have found for the first time that high pressure can be used to make a unique hydrogen-storage material. The discovery paves the way for a new approach to the hydrogen-storage problem. The researchers found that the normally nonreactive, noble gas xenon combines with molecular hydrogen under pressure to form a previously unknown solid with unusual bonding chemistry. The discovery debuts a new family of materials, which could boost hydrogen technologies.

Share:

RELATED TOPICS

Matter & Energy

- Alternative Fuels
- Chemistry
- Organic Chemistry

Earth & Climate

FULL STORY

Related Stories

A New Solution for Storing Energy

May 21, 2014 — Turning concept into a reality, eve bumpy road, but scientist store hydrogen to smooth

Hydrogen R Quickly Whe Nanoparticle

Sep. 29, 2011 — Researc the size of a metal alloy n speed with which hydroge in a metal hydride. The sr read more »



New Metal H Insights Into

Sep. 22, 2011 — A new st of heterometallic molecula features point the way to development of lightweigh read more »

SCIENTIFIC METHOD / SCIENCE & EXPLORATION

Strange, unpredictable chemistry at high pressure

Extreme pressure has "a completely different atomic table," vital to understanding space

by John Timmer - Sep 29, 2013 6:15pm EDT

Share Tweet



The Institute for Advanced Study, which has played host to such luminaries as Albert Einstein and Kurt Gödel, is holding a series of talks to celebrate the birthday of another one of its famous faculty: Freeman Dyson. Dyson made important contributions to a huge variety of fields and gave us the concept of the Dyson Sphere. The talks in his honor covered many of the fields that he influenced, and here, we'll describe the talk by chemist Russel Hemley.

Sci-News.com

Top Headlines:
HD 80606b: Weird Gaseous Exoplanet Raises Questions about Origins of hot Jupiters

'Impossible' Sodium Chlorides Challenge Foundation of Chemistry

Dec 20, 2013 by Editors

« PREVIOUS | NEXT »

Published in
Chemistry

Tagged as
Salt
Sodium chloride

Follow
Like 21K
Share
Tweet
Like 934
G+1 75

A team of researchers led by Prof Artem Oganov of Stony Brook University has shown that, under certain conditions, ordinary rock salt can take on some surprising forms that violate textbook rules of chemistry.



You Might Like



Ice XVII: Chemists Predict Existence of New Form of Ice

LATEST NEWS

Paleontologists Explain Evolution of Sauropod Dinosaurs
Mar 30, 2016 | Paleontology

Moorfields Acuity Chart: Novel Sight Test Detects Early Age-Related Macular Degeneration
Mar 29, 2016 | Medicine

Planetary Researchers Explain Sustained Eruptions on Enceladus
Mar 29, 2016 | Space Exploration

Scientists Produce Map of Neanderthal, Denisovan Ancestry in Present-Day Humans
Mar 29, 2016 | Anthropology

HD 80606b: Weird Gaseous Exoplanet Raises Questions about Origins of hot Jupiters
Mar 29, 2016 | Astronomy

Scientists Produce First Roadmap for European Astrobiology
Mar 28, 2016 | Space Exploration

Scientists Say 'Not Face' is Universal Part of Language
Mar 28, 2016 | Anthropology

Brown Skuas Can Recognize Individual Humans, New Study Shows
Mar 28, 2016 | Biology

NASA's Hubble Space Telescope

PRESSURE

PRESSURE



Under Pressure

- **Orbital hybridization (*s* → *d*)**
- **Complex structures/electronic structure**
- ***Valence-core hybridization at more extreme pressures?***

valence electrons

Static high pressures: Laser-heated DAC







DAC research requires **micro-probing technologies** such as a laser and an intense μm-size x-ray beam at third-generation synchrotron sources

Dynamic High Pressure – Real Extreme Conditions

Gas gun : 300 GPa, 10^3 K



Z pinch : 100 GPa, 10^6 K





core electrons

Tools of research are major facilities

- HPCAT, GSECARS, PETRA, SNS (Current)
- SLAC, DCS-CAT, NIF (Future)

1. CDAC beamtime scheduled every run cycle.
2. Beamtime split between LH and BM-D to allow broader access amongst CDAC partners to the more sought after LH beamtime.
3. Beamtime available for spectroscopy (XANES, EXAFS, IXS) and BM-B based PE cell measurements is becoming crucial for some studies.

CDAC Science

- Discover new Materials (energetic materials, structural materials)
- Understand strategic, superhard Materials – PVT EOS, elastic-plastic deformation, dynamic loading, bonding changes
- Validation of modelling of strategic materials
- Enable the scientific research across all CDAC partners (gas loading, novel cell assemblies, national and international facilties

Novel Xenon Chemistry

JACS
JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

Subscriber access provided by CARNEGIE INST OF WASHINGTON

The xenon-chlorine conundrum: van der Waals complex or linear molecule?

Davide M. Proserpio, Roald Hoffmann, and Kenneth C. Janda

J. Am. Chem. Soc., **1991**, 113 (19), 7184-7189• DOI: 10.1021/ja00019a014 • Publication Date (Web): 01 May 2002




4b1
LUMO
σu
5a1
HOMO
3b1
2b2
Px Py Pz
1a2
2b1
πg
1b2
4a1
πu
3a1
σg
z
y
x
Xe
complex B
Cl–Cl
3σu
LUMO
HOMO
2πu
Px Py Pz
3σg
1πg
1πu
2σu
σu
πu
πg
σg
z
y
x
Xe
Cl–Xe–Cl
Cl···· Cl

Xenon – Chlorine system in a diamond cell

- Mixtures of Xe and Cl_2 compressed in a diamond cell to pressures of the order of 100 GPa display formation of vdW compound at low pressures (2 GPa) that dissociate above 60 GPa (Xe and Cl_2 have similar vdW radii and therefore not surprisingly, we form an fcc (Xe)(Cl_2) compound)

- Enormous increase in intensity of Cl-Cl stretch above 10 GPa (in the Raman) is accompanied by a change in the Xe XANES spectra indicative of change from s to p type character of xenon (Xe k edge is at 34.56 keV)







filled symbols $XeCl_2$ compound;
open symbols pure Cl_2



Orthorhombic $XeCl_2$

Novel Hydride Chemistry

VOLUME 92, NUMBER 18 PHYSICAL REVIEW LETTERS week ending 7 MAY 2004

Hydrogen Dominant Metallic Alloys: High Temperature Superconductors?

N.W. Ashcroft

Laboratory of Atomic and Solid State Physics, Cornell University, Ithaca, New York 14853-2501, USA
Donostia International Physics Center, San Sebastian, Spain

(Received 29 December 2003; published 6 May 2004)

The arguments suggesting that metallic hydrogen, either as a monatomic or paired metal, should be a candidate for high temperature superconductivity are shown to apply with comparable weight to alloys of metallic hydrogen where hydrogen is a dominant constituent, for example, in the dense group IVa hydrides. The attainment of metallic states should be well within current capabilities of diamond anvil cells, but at pressures considerably lower than may be necessary for hydrogen.

PRL **107**, 255503 (2011) PHYSICAL REVIEW LETTERS week ending 16 DECEMBER 2011

Novel Cooperative Interactions and Structural Ordering in H_2S-H_2

Timothy A. Strobel,[1,*] P. Ganesh,[2,†] Maddury Somayazulu,[1] P. R. C. Kent,[2] and Russell J. Hemley[1]







Formation of $Xe(H_2)_8$





Xenon *'dimers'*



electron density of the *vdW compound*







Single Crystal XRD

Unexpected High Pressure Stability of $Xe(H_2)_7$













The ‘dimers’ would reach calculated Xe-Xe distance in hcp xenon at about 110 GPa, but obviously, there is no metallization of the Xe sublattice.

Unexpected Low Pressure Stability of $Xe(H_2)_8$



- $Xe(H_2)_8$ can be synthesized from a mixture of Xe and H_2 at 1.2 GPa and 150 K or 4.2 GPa and 300 K.
- Variable H_2 stoichiometry from a mixture of ordered and disordered H_2 molecules proposed.
- This phase can be retrieved to ambient pressure and temperatures as high as 100 K when H_2 release occurs in a single step.











Novel Energetic Materials

Phys. Rev. B (2015)

Stable Xenon Nitride at High Pressures

Feng Peng[1,2], Yanchao Wang[1], Hui Wang[1], Yunwei Zhang[1], and Yanming Ma[1]*



Figure 2. Crystal structure and bonding properties of XeN_6. (a) XeN_6 in an *R*-3*m* structure without showing any Xe-N bonds at 150 GPa where chaired N_6 hexagons are observed. The small green and large purple balls represent N and Xe atoms, respectively. (b) ELF plots at 150 GPa with an isosurface value of 0.83, showing covalent Xe-N bonding polarized towards N. (c) and (d) depict two structural units of 4-fold bonded N (i.e., N in sp^3 hybridization) and 12-fold bonded Xe, respectively.



Figure 5. Phase diagram of Xe-N_2 system. The dashed line separates the Xe-N_2 mixture (left and green region) and the stable XeN_6 compound (right and blue region).

Xenon – Nitrogen system

- vdW compound identified as $Xe(N_2)_4$ synthesized at 9 GPa and at RT.
- $Xe(N_2)_4$ is stable even at 150 GPa. Turns dark and becomes completely opaque above 120 GPa.
- Xe_xN_y forms when this compound is heated to T > 1500 K for all pressures above 110 GPa



Empty symbols – pure N_2; filled symbols – $Xe(N_2)_4$







High Pressure Alkali Chemistry

PRL **94,** 185502 (2005) PHYSICAL REVIEW LETTERS week ending 13 MAY 2005

Melting of Dense Sodium

Eugene Gregoryanz,[1] Olga Degtyareva,[1] Maddury Somayazulu,[2] Russell J. Hemley,[1] and Ho-kwang Mao[1]



From Metals to Insulators and Back

Formation of Quasimolecules in Insulating States

Interstitial quasiatoms and quasimolecules form when the valence *p* electrons move down relative to s electrons: Li, Na, Mg, Al…



Two nearest ISQ's in *Aba*2-Li form a "molecule"within the structure



Like two H atoms, two ISQs can form both bonding and antibonding orbitals

Charge density for bonding orbital



Charge density for antibonding orbital



The "molecular" orbitals can be understood as maximally localized Wannier functions

isosurface=±2.5

[Miao, Naumov, Hoffmann, & Hemley, *submitted*]

Unexpected Stable Stoichiometries of Sodium Chlorides

20 DECEMBER 2013 VOL 342 SCIENCE

Weiwei Zhang,[1,2]*† Artem R. Oganov,[2,3,4]*† Alexander F. Goncharov,[5,6] Qiang Zhu,[2] Salah Eddine Boulfelfel,[2] Andriy O. Lyakhov,[2] Elissaios Stavrou,[5] Maddury Somayazulu,[5] Vitali B. Prakapenka,[7] Zuzana Konôpková[8]






A
NaCl3+Cl2
60 GPa
Experiment
Le Bail Fit
Experiment x10
Le Bail Fit x10
NaCl3
Cl2
Intensity (arb. units)
Diffraction angle 2θ (degrees)
B
Na3Cl+NaCl
60 GPa
Experiment
Le Bail Fit
Na3Cl P4/mmm
NaCl BCC
Intensity (arb. units)
Diffraction angle 2θ (degrees)
C
Theoretical Pm3n NaCl3
Experimental Pm3n NaCl3
Theoretical Pnma NaCl3
Experimental Pnma NaCl3
Experimental EOS of NaCl B2
Pressure (GPa)
Volume per atom (Å3)
D
Theoretical P4/mmm-Na3Cl
Experimental P4/mmm-Na3Cl
Experimental EOS of NaCl B2
Pressure (GPa)
Volume per atom (Å3)